

The distribution of meadow and steppe vipers (*Vipera graeca*, *V. renardi* and *V. ursinii*): a revision of the New Atlas of Amphibians and Reptiles of Europe

Edvárd Mizsei^{1,2*}, Oleksandr Zinenko³, Neftalí Sillero⁴, Vincenzo Ferri⁵, Stephanos A. Roussos⁶, Márton Szabolcs²

¹Department of Ecology, University of Debrecen, Egyetem tér 1, 4026, Debrecen, Hungary.

²Department of Tisza River Research, Danube Research Institute, Centre for Ecological Research, Hungarian Academy of Sciences, Bem tér 18/c, 4026 Debrecen, Hungary.

³The Museum of Nature; Department of Zoology and Animal Ecology, Biological Faculty, V. N. Karazin Kharkiv National University, Svobody 4, 61022, Kharkiv, Ukraine.

⁴CICGE Centro de Investigação em Ciências Geo-Espaciais, Faculdade de Ciências da Universidade do Porto, Observatório Astronómico Prof. Manuel de Barros, Alameda do Monte da Virgem, 4430-146 Vila Nova de Gaia, Portugal.

⁵Laboratory of Zoology and Evolutionary Biology, University of Rome "Tor Vergata", via Cracovia 1/3, I-00139 Rome, Italy.

⁶Department of Biological Sciences, University of North Texas, Denton, Texas, USA 76203.

*Correspondence: E-mail: edvardmizsei@gmail.com.

Received: 28 November 2017; returned for review: 15 January 2018; accepted 14 March 2018.

In some cases, species' distributions have not been accurate and some geographical areas were not accounted for in the New Atlas of Amphibians and Reptiles of Europe (NA2RE), a project performed by the Mapping Committee of the Societas Europaea Herpetologica (SEH). The distribution of species within the *Vipera ursinii-renardi* group were presented jointly as a single taxon ("*Vipera ursinii/renardi*"), without differentiation between the meadow viper *V. ursinii*, the steppe viper *V. renardi*, and the recently described Greek meadow viper *Vipera graeca*. Here we present a revised 50×50 km resolution distribution map of this group with new records, eliminating putative or erroneous records and extinct populations. Additionally, we filled several spatial gaps in the distribution of *V. renardi* in Eastern Europe by incorporating recent regional overviews, previously considered ambiguous or inaccessible to most researchers, due to language barriers.

Key words: Biogeography, mapping, NA2RE, snake, threatened species, Viperidae.

The New Atlas of Amphibians and Reptiles of Europe (hereafter NA2RE; SILLERO *et al.*, 2014a,b) was created to present and provide accurate knowledge on the spatial distribution of amphibians and reptiles in Europe. It is the most comprehensive and up-to-date spatial database of European herpetofauna. However, exact distributions of some species groups re-

main unresolved (but see WIELSTRA *et al.*, 2014). Species groups have often been the subject of continuous taxonomic studies and rearrangements; thus, species status of some lineages, borders of area, extent of overlaps, and potential hybrid zones are often unknown or unrecognised (e.g. GVOŽDÍK *et al.*, 2010).

The distribution of viper species of the

ursinii-renardi group (sometimes referred as the *Acridophaga* subgenus), have been jointly presented as a single taxon in NA2RE “*Vipera ursinii/renardi*”, including records of both the meadow viper *V. ursinii* and steppe viper *V. renardi*. The latter was not accepted previously as a full species by the Societas Europaea Herpetologica (SEH) (footnote 9 on table 3 in SILLERO *et al.*, 2014a), despite that species delimitation between *V. ursinii* and *V. renardi* had been proved or discerned (NILSON & ANDRÉN, 2001; FERCHAUD *et al.*, 2012; GVOŽDÍK *et al.*, 2012). Further mitochondrial and nuclear DNA evidences also support the species status of these lineages (ZINENKO *et al.* 2015; MIZSEI *et al.*, 2017). A recent taxonomic change has also influenced the current known distribution of this group, as the Greek meadow viper *Vipera graeca*, a former subspecies of *V. ursinii*, was elevated to the species level (MIZSEI *et al.*, 2017). Additionally, most of the distribution within the former Soviet Union is poorly understood because of the lack of accessible observation records. In fact, *V. renardi*, which is more common in lowland steppes, is more widespread than previously depicted in the NA2RE (e.g. ZINENKO *et al.* 2015). Most published occurrence records are hard to find or overlooked, as they are often harboured in regional publications and written in local languages. Here, we have compiled these observations to update the map.

We revised the presence records of the group, differentiating *V. ursinii*, *V. renardi* and *V. graeca* in the NA2RE data (Supplementary Material) cell by cell. We have (i) identified cells with species' presence (=1); (ii) confirmed species' presence

with available data in published or unpublished sources; (iii) added new species' presence when we had available data for a cell previously showing species' absence (=0); (iv) deleted species' presence (1→0) when there was no evidence of presence (mainly by erroneous identification of species and locations); (v) changed status of cells to historical where there was no observations for the last 25 years, and provided the date of last observations; and finally (vi) we added a comment to each cell where changes were implemented. We determined the species according to recent molecular studies on meadow and steppe vipers (FERCHAUD *et al.*, 2012; ZINENKO *et al.*, 2015; MIZSEI *et al.*, 2017). Following the results of FERCHAUD *et al.* (2012) and ZINENKO *et al.* (2015), we treated *Vipera lotievi* (NILSON *et al.*, 1995) as *V. renardi* s.l. To revise the distribution we used literature sources (SUKHOV, 1928; BANNIKOV *et al.*, 1977; NILSON & ANDRÉN, 2001; VLASOV & VLASOVA, 2001; DOTSSENKO, 2003; KRECSÁK *et al.*, 2003; FILIPPI & LUISELLI, 2004; FERRI & MARCONI, 2006; GHIRA, 2007; VEDMEDERJA *et al.*, 2007; ZINENKO & BAKIEV, 2007; FERRI & PELLEGRINI, 2008; SELIUNINA, 2008; KOTENKO & KUKUSHKIN, 2008; 2009; STRUGARIU *et al.*, 2011; ZAMFIRESCU *et al.*, 2012; COGĂLNICEANU *et al.*, 2013; DEBELO & CHIBILYOV, 2013; FROLOVA & KLIMOV, 2013; JELIĆ *et al.*, 2013; LYET *et al.*, 2013; BAKIEV *et al.*, 2015; PÉCHY *et al.*, 2015; TUPIKOV & ZINENKO, 2015; ZINENKO *et al.*, 2015; BAKIEV *et al.*, 2016; MIZSEI *et al.*, 2016); museum collections (The M. Shcherbak Zoological Museum, National Museum of Natural History at the National Academy of Sciences of Ukraine, Kiev; The Museum of Nature at V. N. Karazin Kharkiv National Universi-




Figure 1: Revised distribution of meadow and steppe vipers in Europe.

ty, Kharkiv, Ukraine); and records from our personal unpublished databases, and other unpublished sources (see acknowledgements). The date of each record was taken from the corresponding publication; in the case when no date was available in the text, we used the date of the publication for which it was referenced in. All records before 1992 (from 25 years ago) were considered as “historical”. After that point, with the maximum pressure from

agriculture on steppic habitats of *V. renardi* and *V. ursinii* in the former USSR countries, there was temporary relief for steppic habitats, and a shift in the way of their management. Simultaneously, national Red data books and sometimes local monitoring programs started to gather current data about local species in newly established independent countries.

We added seven cells to the distribution of *V. graeca* from new populations recent-

ly discovered (Fig. 1). Furthermore, only one cell was confirmed and another one was deleted because observations suggest presence only in the neighbouring cell. We validated the presence of *V. renardi* in 42 cells, added 173 new cells based on published (N = 144 cells), and unpublished observations of the authors (N = 29 cells), listed 69 as historical, and deleted 6 cells (erroneous species identification in the literature; specification and correction of geographic coordinates of localities), most of them located in the north-eastern edge of the species' distribution in Europe. However, we were unable to validate 56 cells where *V. renardi* is present in the NA2RE, but kept these cells as occupied in the database until these areas are further investigated. The distribution of *V. ursinii* (separated from the distribution of *V. graeca* and *V. renardi*) remained the same in most parts (33 cells), but we added 17 cells mainly in the Balkan Peninsula based on the comprehensive database of JELIĆ *et al.* (2013). We have changed the status of 17 cells to "historical", mostly in Bessarabia (Ukraine, Romania, and Republic of Moldova), the eastern edge of the species range (*V. ursinii* ssp. *moldavica*). Most likely, all populations have gone extinct in these areas due to habitat alteration by agriculture. Last recorded observations of meadow viper there belong to the 1970's, and only a few survived populations near Iasi and in the Danube Delta still existing (KRECSÁK *et al.*, 2003). There are no known overlaps among the geographic ranges of the three species, thus, we consider them as allopatric; however, a contact zone may have historically existed between *V. renardi* and *V. u. moldavica* (see Fig. 1). Overall,

we increased the known distribution of *V. graeca* by 87.5%, *V. renardi* by 63.8%, and *V. ursinii* by 34.0% in 50×50 km grid cells.

We assume that at this scale (50×50 km grids), further fieldwork may only slightly increase the currently known range of *V. graeca* and *V. ursinii*. However, it is possible to improve our knowledge of the distribution at finer scales, mostly in the reticulating mountain ranges in the Balkan Peninsula where these species live in isolated high alpine meadows that have been poorly explored. Due to the large distribution of *V. renardi* in Eastern Europe, which was mostly continuous in the past, there is potential for substantially extending the list of localities of these species and filling gaps in the known distribution, mostly in Southern Russia. Greater prevalence of "historical" observations (before 1992) in that region, indicates a need for contemporary surveys and consistent monitoring. Recent habitat loss, population size decrease, and population extinctions may be overlooked due to data deficiency. Northern isolated historical records in Russia may have resulted from range fluctuations in postglacial times (ZINENKO & BAKIEV, 2007), similarly to other reptile species in the region (MAROSI *et al.*, 2012); many of these populations are most likely to be extinct now.

All three species are threatened by past and ongoing habitat loss and fragmentation. IUCN Red List Category of *V. ursinii* is Vulnerable (VU) globally (JÖGER *et al.*, 2009), and *V. renardi* is VU in Europe regionally (COX & TEMPLE, 2009). *V. graeca* fulfils the criteria for the category of Endangered (EN) because the species' area of occupancy is less than 500 km², the distribution is severely fragmented and there is ongoing habitat loss

and degradation (MIZSEI *et al.* in press). Our updated maps increase vital information needed for the implementation of effective conservation of these species, by pinpointing the geographic areas to focus future conservation efforts on. Our results are integrated to NA2RE, the web-based spatial data infrastructure, underpinning the New Atlas of Amphibians and Reptiles of Europe (<http://na2re.ismai.pt>).

Acknowledgement

EM and MS were granted with the Societas Europaea Herpetologica, SEH Grant in Herpetology, Mapping and Distributions in 2015 and EM, SAR, MS and OZ in 2016. MS was supported by Balassi Institute Campus Hungary Programme (B2/1CS/11521) and OTKA (K106133). EM and MS were granted by Balassi Institute Campus Hungary Programme (B2/1CS/19196). NS is supported by an IF contract (IF/01526/2013) by Fundação para a Ciência e Tecnologia (FCT, Portugal). OZ is supported within the project “Biological challenges and threats due to migrations and invasions” funded by Ukrainian ministry of science and education; and a Rufford Small Grant (14742-1). Finally, we thank to the three anonymous reviewers to their comments to the earlier version of the manuscript. We are grateful to G. Păun for providing unpublished data.

REFERENCES

- BAKIEV, A.G.; GORELOV, R.A.; KLYONINA, A.A.; RYZHOV, M.K. & SOLOMAYKIN, E.I. (2016). Snakes from the Red book of the Samara area: new places of finds. *Samarskaya Luka: problems of regional and global ecology* 25: 129-130.
- BAKIEV, A.G.; GARANIN, V.I.; GELASHVILI, D.B.; GORELOV, R.A.; DORONIN, I.V.; ZAITSEVA, O.V.; ZINENKO, O.I.; KLYONINA, A.A.; MAKAROVA, T.N.; MALENYOV, A.L.; PAVLOV, A.V.; PETROVA, I.V.; RATNIKOV, V.YU.; STARKOV, V.G.; SHIRYAEVA, I.V.; YUSUPOV, R.KH. & YAKOVLEVA, T.I. (2015). *The Vipers (Reptilia: Serpentes: Viperidae: Vipera) of the Volga basin*. Part 1. Tolyatti, Kassandra. [in Russian]
- BANNIKOV, A.G.; DAREVSKY, I.S.; ISHCENKO, V.G.; RUSTAMOV, A.K. & SHCHERBAK, N.N. (1977). *Guide to Amphibians and Reptiles of the USSR*. Moscow, Prosveshchenie. [in Russian]
- COGĂLNICEANU, D.; ROZYLOWICZ, L.; SZÉKELY, P.; SAMOILĂ, C.; STĂNESCU, F.; TUDOR, M.; SZÉKELY, D. & IOSIF, R. (2013). Diversity and distribution of reptiles in Romania. *ZooKeys* 341: 49-76.
- COX, N.A. & TEMPLE, H.J. (2009). *European Red List of Reptiles*. Luxembourg, Office for Official Publications of the European Communities.
- DEBELO, P.V. & CHIBILYOV, A.A. (2013). *The Amphibians and Reptiles of the Ural-Caspian region. Natural diversity in the Ural - Caspian region series. Vol. 3*. Yekaterinburg, Ural branch of Russian Academy of Sciences.
- DOTSENKO, I.B. (2003). *Catalogue of the collections of Zoological Museum, NSNHM, NAS of Ukraine*. The snakes. Kiev, NSNHM.
- FERCHAUD, A.-L.; URSENBACHER, S.; CHEYLAN, M.; LUISELLI, L.; JELIĆ, D.; HALPERN, B.; MAJOR, Á.; KOTENKO, T.; KEYAN, N.; CRNOBRNJAIŠAILOVIĆ, J.; TOMOVIĆ, L.; GHIRA, I.; IOANNIDIS, Y.; ARNAL, V. & MONTGERALD, C. (2012). Phylogeography of the *Vipera ursinii* complex (Viperidae): mitochondrial markers reveal an east-west disjunction in the Palearctic region. *Journal of Biogeography* 39: 1836-1847.
- FERRI, V. & MARCONI, M. (2006). *Vipera di Orsini, Vipera ursinii* (Bonaparte, 1835), In R. SINDACO; G., DORIA, E., RAZZETTI & F., BERNINI (eds.) *Atlante degli anfibi e dei rettili d'Italia / Atlas of Italian Amphibians and Reptiles*. Societas Herpetologica Italica, Edizioni Po-

- listampa, Firenze, pp. 606-611.
- FERRI, V. & PELLEGRINI, MR. (2008). *Vipera* di Orsini, *Vipera ursinii* (Bonaparte, 1835), In L., DI TIZIO; MR., PELLEGRINI; N., DI FRANCESCO & M., CARAFA (eds.) *Atlante dei Rettili d'Abruzzo*. Ianieri-Talea Edizioni, Pescara, pp. 184-189.
- FILIPPI, E. & LUISELLI, L. (2004). Ecology and conservation of the Meadow viper, *Vipera ursinii*, in three protected mountainous areas in central Italy. *Italian Journal of Zoology* 71: 159-161.
- FROLOVA, E.N. & KLIMOV, A.S. (2013). Some morphological characters of the steppe viper (*Pelias renardi*) vicinity of museum-reserve "Divnogorye". *Modern herpetology: problems and ways of their solutions*. The first international conference of the young herpetologists of Russia and neighbouring countries. Saint-Petersburg.
- GHIRA, I. (2007). Rediscovery of *Vipera ursinii rakosiensis* in Transylvania. *Herpetologica Romanica* 1: 77-81.
- GVOŽDÍK, V.; JANDZIK, D.; CORDOS, B.; REHÁK, I. & KOTLÍK, P. (2012). A mitochondrial DNA phylogeny of the endangered vipers of the *Vipera ursinii* complex. *Molecular Phylogenetics and Evolution* 62: 1019-1024.
- GVOŽDÍK, V.; JANDZIK, D.; LYMBERAKIS, P.; JABLONSKI, D. & MORAVEC, J. (2010). Slow worm, *Anguis fragilis* (Reptilia: Anguidae) as a species complex: Genetic structure reveals deep divergences. *Molecular Phylogenetics and Evolution* 55: 460-472.
- JELIĆ, D.; AJTIĆ, R.; STERIJOVSKI, B.; CRNOBRNJA-ISAILOVIĆ, J.; LELO, S. & TOMOVIĆ, L. (2013). Distribution of the genus *Vipera* in the western and central Balkans (Squamata: Serpentes: Viperidae). *Herpetozoa* 25: 109-132.
- JÖGER, U.; CRNOBRNJA-ISAILOVIĆ, J.; VOGGRIN, M.; CORTI, C.; STERIJOVSKI, B.; WESTERSTRÖM, A.; KRECSÁK, L.; PÉREZ MELLADO, V.; SÁ-SOUSA, P.; CHEYLAN, M.; PLEGUEZUELOS, J. M. & SINDACO, R. (2009). *Vipera ursinii*. *The IUCN Red List of Threatened Species* 2009: e.T22997A9406628. Retrieved on 11/05/2016.
- KOTENKO, T.I. & KUKUSHKIN, O.V. (2009). Steppe viper, *Vipera renardi* (Christ.), In I. A., AKIMOV (ed.) *The Red Data Book of Ukraine. Animals*, Kiev, Globalkonsalting.
- KOTENKO, T.I. & KUKUSHKIN, O.V. (2008). Steppe viper, *Vipera renardi* (Christ.), - a species of the Red Data Book of Ukraine. Kiev, *Registration of Animals under Red Data Book of Ukraine*.
- KRECSÁK L.; ZAMFIRESCU S. & KORSÓS Z. (2003): An updated overview of the distribution of the Moldavian steppe viper (*Vipera ursinii moldavica* Nilson, Andren et Joger, 1993). *Russian Journal of Herpetology* 10: 199-206.
- LYET, A.; THUILLER, W.; CHEYLAN, M. & BESNARD, A. (2013). Fine-scale regional distribution modelling of rare and threatened species: bridging GIS Tools and conservation in practice. *Diversity and Distribution* 19: 651-663.
- MAROSI, B.; ZINENKO, O.I, GHIRA, I.V.; CRNOBRNJA-ISAILOVIC, J.; LYMBERAKIS, P.; SOS, T. & POPESCU, O. (2012). Molecular data confirm recent fluctuations of northern border of dice snake (*Natrix tessellata*) range in Eastern Europe. *North-Western Journal of Zoology* 8: 374-377.
- Mizsei, E., Szabolcs, M., Dimaki, M., Roussos, S.A., Ioannidis, Y. (in press): *Vipera graeca*. The IUCN Red List of Threatened Species.
- MIZSEI, E.; JABLONSKI, D.; ROUSSOS, S.A.; DIMAKI, M.; IOANNIDIS, Y.; NILSON, G. & NAGY, Z.T. (2017). Nuclear markers support the mitochondrial phylogeny of *Vipera ursinii-renardi* complex (Squamata: Viperidae) and species status for the Greek meadow viper. *Zootaxa* 4227: 75-88.
- MIZSEI, E.; ÜVEGES, B.; VÁGI, B.; SZABOLCS, M.; LENGYEL, S.; PFLIEGLER, W.P.; NAGY, Z.T. & TÓTH, J.P. (2016). Species distribution modelling leads to the discovery of new populations of one of the least known European snakes, *Vipera ursinii graeca*, in Albania. *Amphibia-Reptilia* 37: 55-68.
- NILSON, G. & ANDRÉN, C. (2001). The Meadow and Steppe Vipers of Europe and Asia - The Vi-

SHORT NOTES

- pera* (Acridophaga) ursinii complex. *Acta Zoologica Academiae Scientiarum Hungaricae* 47: 87-267.
- NILSON, G.; TUNIYEV, B. S.; ANDRÉN, C. & ORLOV, N. (1995). Systematics of the vipers of the Caucasus: Polymorphism or sibling species? *Asiatic Herpetological Research* 6: 1-16.
- PÉCHY, T.; HALPERN, B.; SÓS, E. & WALZER, C. (2015). Conservation of the Hungarian meadow viper *Vipera ursinii rakosiensis*. *International Zoo Yearbook* 49: 89-103.
- SELIUNINA, Z.V. (2008). Reptiles of the Black sea biosphere reserve in 1990-2005. Kiev, *Registration of Animals under Red Data Book of Ukraine*.
- SILLERO, N.; CAMPOS, J.; BONARDI, A.; CORTI, C.; CREEMERS, R.; CROCHET, P.-A.; CRNOBRNJA ISAILOVIĆ, J.; DENOËL, M.; FICETOLA, G.F.; GONÇALVES, J.; KUZMIN, S.; LYMBERAKIS, P.; DE POUS, P.; RODRÍGUEZ, A.; SINDACO, R.; SPEYBROECK, J.; TOXOPEUS, B.; VIEITES, D.R. & VENCES, M. (2014a). Updated distribution and biogeography of amphibians and reptiles of Europe. *Amphibia-Reptilia* 35: 1-31.
- SILLERO, N.; OLIVEIRA, M.A.; SOUSA, P.; SOUSA, F. & GONÇALVES-SECO, L. (2014b). Distributed database system of the New Atlas of Amphibians and Reptiles in Europe: the NA2RE project. *Amphibia-Reptilia* 35: 33-39.
- STRUGARIU, A.; ZAMFIRESCU, S.R.; GHERGHEL, I.; SAHLEAN, T.C.; MORARU, V. & ZAMFIRESCU, O. (2011). A preliminary study on population characteristics and ecology of the critically endangered meadow viper *Vipera ursinii* in the Romanian Danube Delta. *Biologia* 66: 175-180.
- SUKHOV, G.F. (1928). Materials to the herpetofauna of Poltava region. *Proceedings of Poltava state Museum* 1: 251-256.
- TUPIKOV, A.I. & ZINENKO, O.I. (2015). Distribution of the steppe viper *Vipera renardi* (Reptilia, Viperidae) in Kharkiv region. *Visnyk of Dnipropetrovsk University. Biology, Ecology* 23: 172-176.
- VEDMEDERJA, V. I.; ZINENKO, O.I. & GONCHARENKO, L. A. (2007). *Catalogue of collections of the Museum of Nature at V. N. Karazin's Kharkiv National University*. Snakes (Reptilia: Serpentes). Kharkiv.
- VLASOV, A.A. & VLASOVA, O.P. (2001). *Reintroduction of the steppe viper (Vipera ursinii renardi) in forest-steppe nature reserves of the Central Chernozemje*. Questions of herpetology. Proceedings of the First meeting of A. M. Nikolsky Herpetological Society. pp. 57-58. Moscow, Puschino.
- WIELSTRA, B.; SILLERO, N.; VÖRÖS, J. & ARNTZEN, W. (2014). The distribution of the crested and marbled newt species (Amphibia: Salamandridae: *Triturus*) – an addition to the New Atlas of Amphibians and Reptiles of Europe. *Amphibia-Reptilia* 35: 371-381.
- ZAMFIRESCU, S. R.; STRUGARIU, A.; ZAMFIRESCU, O. & GHERGHEL, I. (2012). In situ confirmation of the occurrence of the critically endangered Moldavian meadow viper (*Vipera ursinii moldavica*) in the Cîrteii Valley (Iasi county, Romania). *North-Western Journal of Zoology* 8: 378-381.
- ZINENKO, O.I. & BAKIEV, A.G. (2007). About Change of the Northern Boundary of Distribution of the Steppe Viper, *Vipera renardi* (Reptilia, Viperidae), in European Russia. *Vestnik Zoologii* 41: 478.
- ZINENKO, O.; STÜMPPEL, N.; MAZANAIEVA, L.; BAKIEV, A.; SHIRYAEV, K.; PAVLOV, A.; KOTENKO, T.; KUKUSHKIN, O.; CHIKIN, Y.; DUSEBAYEVA, T.; NILSON, G.; ORLOV, N.L.; TUNIYEV, S.; ANANJEVA, N.B.; MURPHY, R.W. & JOGER, U. (2015). Mitochondrial phylogeny shows multiple independent ecological transitions and northern dispersion despite of Pleistocene glaciations in meadow and steppe vipers (*Vipera ursinii* and *Vipera renardi*). *Molecular Phylogenetics and Evolution* 84: 85-100.